

Role of Ultrasonography in Thyroid Nodules with Pathological Correlation

Pranay Kumar¹, Subhash C. Sylvania², Abhay Kumar Singh³

¹Post Graduate IIIrd year, ²Professor, Department of Radiodiagnosis, ³Assistant Professor, Department of ENT, Saraswathi Institute of Medical Sciences Hapur (U.P)

Abstract

Introduction: Thyroid nodule imaging had been a challenging task. Ultrasound has been the modality of choice for detection & evaluation of these lesions. This study was done to identify the accuracy of various imaging features in thyroid nodules that are associated with benignity and malignancy and the overall accuracy of ultrasound in determining malignant nodules.

Objective: Ultrasonographic evaluation of thyroid to characterize the thyroid disease and nodules (benign vs malignant) and their morphologic evaluation .

Materials and Method: A total of Seventy patients (70) of both sexes and different age group, meeting the inclusion criterion, were the part of present prospective study. The cases referred to the Department of Radio-diagnosis and imaging from OPD/IPD of Saraswathi Institute of Medical sciences, Hapur from October'2017 to July'2019. An informed consent was taken from all the patients subjected for evaluation.

Results: The collective data was analysed for Ultrasound findings of thyroid nodule which presented clinically as solitary or multiple thyroid nodules. The total number of cases analysed was 70 out of which 48 were found to have solitary/multinodular lesions clinically. On ultrasound evaluation out of 48 patients 38 had solitary/multinodular lesions.

Conclusion: High resolution sonography is recommended as the primary imaging modality in evaluation of thyroid diseases. It has a high sensitivity and specificity in the diagnosis of thyroid diseases.

Keywords: High resolution ultrasonography, multinodular, thyroid

Introduction

Thyroid nodules are extremely common, found at palpation in 4% to 7% of an asymptomatic population.^[2] in 17% to 27% of cases at USG.^[3-5] and in 50% of cases at autopsy.^[6] USG is the modality of choice in a clinically suspected case of thyroid lesion and the most sensitive imaging modality to detect a clinically non-palpable nodule of 2-3 mm size and allows a more accurate

morphological characterization of the lesion including the size and the number of thyroid nodules, the volume of thyroid tissue and could well differentiate with extra-thyroid neck masses. An addition of color and spectral Doppler imaging that determines the vascular pattern of thyroid diseases has been found to be very useful tool in screening the thyroid nodule for malignancy. ^[1]

Material and Method

This prospective study was conducted from 1st October 2017 to 30th July' 2019 on the cases referred from OPD/IPD of the SIMS Hospital, to the Department of Radio-diagnosis, SIMS, Hapur in collaboration with the Department of Pathology, Otolaryngology, and Surgery. Sonography was performed using Philips

Corresponding Author:

Dr. Pranay Kumar

Post Graduate IIIrd year

Department of RADIODIAGNOSIS

Saraswathi Institute of Medical Sciences, Hapur (U.P)

Harypol278@gmail.com, Cont.: 9464780122

clear view 350 USG unit with multi-frequency probe of 5.0 to 13MHz probe. A phased array curvilinear broadband transducer of 2 to 5 MHz frequency was also used to assess larger lesions. Fine needle aspiration cytology(FNAC) and Histo-pathological correlation, if patient is subjected to surgery, in thyroid diseases was also done. Clinical findings or investigations and imaging findings were co – related. Distribution of cases according to post ultrasound diagnosis was done.

Inclusion criteria

- • All the cases presented and referred on clinical grounds for suspicion of thyroid diseases with or without deranged thyroid function test.
- • Patients who presented with a palpable thyroid nodule.
- • Patients with deranged thyroid function tests.
- • Incidentally discovered thyroid nodules.

Exclusion criteria

- • All patients with previous FNAC or biopsy proven diagnosis.
- • Patients who had previously undergone thyroid surgery for any reason.

Results

In the present study, there was an female preponderance and out of the 70 patients 61/70 (87.4%) were females and 9/70 (12.6%) were males. The M:Fratio in the study group is 1: 6.7 showed a female preponderance for thyroid diseases. The study group composed of patients ranging from 7 to 92 years of age. Majority of the patients were in the age group of 31-40years, accounting for 31.5% of the cases. The mean age in the study was 33.6years. Only one patient above 70 years of age was included in the study. Female preponderance was noted in patients with thyroid disorders in all age groups except between 61 to 70 years, where there was equal ratio of males and females. Five patients with clinical suspicion of thyroid disease, were found normal at sonography.

Out of 48/70 (68.57%) patients presented with solitary thyroid nodule on clinical grounds and out of

these 17/48 (35.4%) of patients diagnosed as Solitary Thyroid Nodule (STN) clinically were found to have a multi-nodular (MNG) thyroid disease at high resolution USG. 21/48(43.75%) patients were found to have sonographic STN while 10/48(20.85%) of these patients had absence of nodules on ultrasound.

Most of the nodules of colloid goiter were isoechoic 16/21(76.4%), followed by heteroechoic in 14.2%. Consistency of 12/21 (57%) nodules was mixed solid- cystic, followed by solid nodules in 28.5%. Only 14.2% of cases were purely cystic in consistency.

Calcification was seen in 9 (42.7%) cases out of 21 nodules, among which 28.5% were coarse calcifications and 14.2% were comet tail type of calcification. A thin well defined, complete halo was seen in 85.7% of these lesions. Incomplete halo was seen 14.3% of cases. Most of the nodules (15/21) showed a predominantly peripheral flow pattern (71.4%). The 23.8% of the nodules showed both peripheral and intranodular vascularity. Only one nodule showed absence of vascularity.

17 out of 70 patients were diagnosed as multinodular goitre (MNG) on ultrasonography and were subsequently proved on FNAC. 70.5% of the MNG were solid- cystic in consistency, followed by purely solid consistency in 23.5% and purely cystic in 6% cases. 52.9% of cases were isoechoic in echopattern, followed by heteroechoic echopattern in 35.5%. Anechoic and hyperechoic were seen in 5.8% of cases (1) each. Calcifications were present in 29.4% (5) cases and all were coarse type. Thin regular complete halo were present in 88.3% (15) cases. Halo was absent in 11.7% (2) cases. On CDI, internodular vascularity was seen in 82.6% cases followed by intranodular flow in 5.8% cases. No vascularity (absent flow) was seen in 11.6% cases.

Two patients of Grave's disease had solid, enlarged, bulky, diffusely hypoechoic thyroid with few echogenic strands. The patients exhibited the thyroid inferno pattern at colour flow imaging.

89.4% of the cases presented with hypoechoic echotexture and 10.6% cases presented with heteroechoic echotexture. There was nodularity in two of the patients in the form of discrete nodules in a background of diffusely altered parenchyma.

Two cases of papillary neoplasm showed heteroechoic thyroid parenchyma with ill defined multiple nodules with solid-cystic changes and cystic degeneration with absence of halo. Both cases showed presence of microcalcifications. Extrathyroid involvement in the form of lymph nodes was seen in both cases.

Discussion

The present study was undertaken to evaluate the role of High Resolution Ultra-Sonography in evaluation of thyroid diseases. The largest group of patients were in the 31-40 year age group with 87.4% of the patients were females and 12.6% were males. The overall sex ratio was M:F = 1:6.7. A similar female preponderance was noted by Solbiati et al^[9] in 1985 and Nam Goong et al^[12] in 2003 in their respective studies.

Out of 48 patients with clinical suspicion of thyroid nodule, 35.4% of patients diagnosed as Solitary Thyroid Nodule clinically were found to have a multinodular thyroid disease at high resolution sonography. 21(43.75%) patients were sonographically proven to be STN. 10 patients who clinically suspected to be STN were diagnosed as other thyroid disorders and showed no evidence of nodularity on Ultra-sonography.

Of the 38 patients with non toxic hyperplastic goiter, 21 patients had solitary thyroid nodule while 17 had multinodular goiter. In patients with STN, consistency of 57% nodules was mixed solid- cystic, followed by solid nodules in 28.5%. Only 14.2% of STN cases were purely cystic in consistency. 70.5% of the MNG were solid- cystic in consistency, followed by purely solid consistency in 23.5% cases.

Thyroid calcifications can be classified as coarse calcification, peripheral or microcalcifications. Thyroid microcalcifications are psammoma bodies which are 10-100um round laminar calcific deposits. At USG, microcalcifications appear as punctate hyperechoic foci without acoustic shadowing. Microcalcifications are found in 29-59% of all thyroid carcinomas^[7,8], most commonly in papillary thyroid carcinomas.

In the present study, both the cases of papillary thyroid carcinoma showed presence of microcalcification. In 2004, Iannucilli et al stated that

presence of microcalcifications is the only statistically significant predictor of malignancy in thyroid nodules.^[15]

Peripheral calcification is seen most commonly in multinodular goitre but may also be present in malignancy^[13]. 5.8% of MNG cases showed presence of peripheral calcification in the present study.

Invasion of anatomic structures around thyroid was observed in 1(50%) patient of malignancy. The patient showed extension of the lesion in the tracheoesophageal groove with infiltration into the trachea which was confirmed on CT. Local invasion into strap muscles, tracheal infiltration and carotid entrapment was seen in 26% of the thyroid malignancies in the study group of Hayashi et al 1986.^[11]

Cervical adenopathy was seen in both cases of thyroid malignancy. One patient showed presence of microcalcification in the lymph nodes as well. Solbiati 1992 described microcalcification and cystic degeneration in the nodes as pathognomic for metastatic nodes from thyroid primary.^[16]

Confirmation of thyroid tissue was done by FNAC and thyroid scintigraphy.

Results of our study show that high resolution sonography had sensitivity of 96.88% and specificity of 83.33% in diagnosing thyroid lesions with a positive predictive value of 98.41% and NPV of 71.43%.

Conclusion

High resolution sonography is a useful modality for evaluation of thyroid diseases. It is reliable in distinguishing normal from abnormal thyroid. Thyroid sonography is useful in defining whether the patient has a diffuse abnormality, a multinodular pathology or a solitary nodule. It is an excellent modality for morphological characterization of thyroid lesions. High resolution sonography can differentiate benign from malignant thyroid nodules and masses in majority of cases. Sonography is very sensitive in detecting local invasion by thyroid malignancies. However, the substernal and retrosternal components of thyroid masses cannot be adequately imaged due to technical constraints. It is useful in diagnosis and follow up of diffuse thyroid diseases i.e. Grave's disease and

thyroiditis. Sonography can easily and reliably exclude the possibility of ectopic thyroid in suspected cases of thyroglossal cysts or midline neck swellings. It is a useful imaging modality in evaluation of thyroid in children and pregnant women. Sonography can be used precisely to guide FNAC from impalpable thyroid lesions. Color flow imaging gives useful information about the vascular status of the lesions in thyroid and helps in characterizing indeterminate lesions in some cases.

Ethical Clearance- Taken from ethical committee of institute

Source of Funding- Self

Conflict of Interest – Nil

References

1. Chaudhary V, Bano S. Imaging of the thyroid: Recent advances. *Indian J EndocrMetab* 2012; 16: 371-6
2. Mazzaferri E. Management of a solitary thyroid nodule. *N Engl J Med* 1993; 328: 553–559.
3. Carroll B. Asymptomatic thyroid nodules: incidental sonographic detection. *AJR Am J Roentgenol* 1982; 138:499–501.
4. Woestyn J, Afschrift M, Schelstraete K, Vermeulen A. Demonstration of nodules in the normal thyroid by echography. *Br J Radiol* 1985; 58:1179–1182.
5. Brander A, Viikinkoski V, Nickels J, Kivisaari L. Importance of thyroid abnormalities detected at US screening: a 5-year follow-up. *Radiology* 2000; 215: 801–806.
6. Mortenson J, Woolner L, Bennett W. Gross and microscopic findings in clinically normal thyroid glands. *J Clin Endocrinol Metab* 1955; 15:1270–1280.
7. Papini E, Guglielmi R, Bianchini A, et al. Risk of malignancy in non-palpable thyroid nodules: predictive value of ultrasound and color Doppler features. *J Clin Endocrinol Metab*, May 2002, 87(5):1941–1946
8. Ayesha T. Khizer, Saleem Raza and Atiq-ur-Rehman Slehria: Diffusion-weighted MR Imaging and ADC Mapping in Differentiating Benign from Malignant Thyroid Nodules. *Journal of the College of Physicians and Surgeons Pakistan* 2015, Vol. 25 (11): 785-788
9. Solbiati L, D Pra L, Lerace T et al. High resolution sonography of the recurrent laryngeal nerve: Anatomic and Pathologic considerations. *AJR* 1985 Nov; 989-993.
10. James EM, Charboneau JW. High frequency thyroid ultrasonography. *Seminar in US, CT and MR* 1985 Sept;6(3) : 294-309
11. Hayashi N, Tamaki N, Yamamoto K et al. Real time ultrasonography of thyroid nodules. *Acta Radiologica Diagnosis* 1986; 27 :404-408.
12. Nam-Goong IS , Kim HY , Gong G , et al . Ultrasonography-guided fine-needle aspiration of thyroid incidentaloma: correlation with pathological findings . *Clin Endocrinol (Oxf)* 2004 ; 60 (1): 21 – 28 .
13. Hayashi N, Tamaki N, Yamamoto K et al. Real time ultrasonography of thyroid nodules. *Acta Radiologica Diagnosis* 1986; 27 :404-408.
14. Nam-Goong IS , Kim HY , Gong G , et al . Ultrasonography-guided fine-needle aspiration of thyroid incidentaloma: correlation with pathological findings . *Clin Endocrinol (Oxf)* 2004 ; 60 (1): 21 – 28 .
15. Kovacevic DO, Skurla MS. Sonographic diagnosis of thyroid nodules: correlation with the results of sonographically guided FNAB. *J Clin ultrasound* 2007 Feb; 35(2): 63-7.
16. Clark KJ, Cronan JJ, Scola FH. Color Doppler Sonography: Anatomic and Physiologic assessment of the thyroid. *J Clin Ultrasound* 1995 May 23; 23: 215-223.
17. Iannuccilli JD, Cronan JJ, Monchik JM. Risk for malignancy of thyroid nodules as assessed by sonographic criteria: the need for biopsy. *J ultrasound Med* 2004; 23(11) : 1455-64.
18. Solbiati L, Cioffi V, Ballarati E. Ultrasonography of neck. *RCNA* 1992 Sept; 30(5): 941-54.
19. Frates MC, Benson CB, Charboneau JW et al., “Management of thyroid nodules detected at US: society of Radiologists in Ultrasound consensus conference statement,” *Ultrasound* 2006; 22: 231–238.